

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁵ : A61K 31/48, 47/36, 47/40</p>	<p>A2</p>	<p>(11) International Publication Number: WO 94/22445 (43) International Publication Date: 13 October 1994 (13.10.94)</p>
<p>(21) International Application Number: PCT/EP94/00891 (22) International Filing Date: 18 March 1994 (18.03.94) (30) Priority Data: 9300297 26 March 1993 (26.03.93) BE 9300298 26 March 1993 (26.03.93) BE 9300299 26 March 1993 (26.03.93) BE (71)(72) Applicant and Inventor: MERKUS, Franciscus, W., H., M. [NL/BE]; Antwerpsesteenweg 165, B-2350 Vosselaar (BE). (74) Agent: LOUËT FEISSER, Arnold; Trenité Van Doorne, P.O. Box 75265, NL-1070 AG Amsterdam (NL).</p>		<p>(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: PHARMACEUTICAL COMPOSITIONS FOR INTRANASAL ADMINISTRATION OF DIHYDROERGOTAMINE, APO-MORPHINE AND MORPHINE (57) Abstract <p>The invention relates to pharmaceutical compositions for the intranasal administration of dihydroergotamine, apomorphine and morphine comprising one of these pharmacologically active ingredients in combination with a cyclodextrin and/or a disaccharide and/or a polysaccharide and/or a sugar alcohol.</p></p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PHARMACEUTICAL COMPOSITIONS FOR INTRANASAL ADMINISTRATION
OF DIHYDROERGOTAMINE, APOMORPHINE AND MORPHINE

- 5 This invention is related to pharmaceutical compositions
for nasal administration of dihydroergotamine, apomorphine
and morphine, and methods of administering such
compositions.
- 10 Dihydroergotamine mesylate (DHE) has been used in migraine
therapy already for a long time. In patients with migraine
attacks, DHE is suitable for basic interval treatment using
tablets or solution, both for oral application, as well as
for acute treatment by intravenous or intramuscular
15 injection. DHE has been introduced in a nasal spray to
avoid the parenteral and the oral route of administration.
The nasal spray seems a good alternative, because it is
less painful, less expensive and less inconvenient than
injection therapy. Secondly, nausea and vomiting are common
20 in migraine patients, making a nasal spray much more
efficient than oral treatment.

- A nasal spray containing DHE 4 mg/ml in an aqueous solution
has been studied extensively by a number of investigators.
25 Some of these investigators report, that besides DHE the
nasal spray also contains glucose 5% and caffeine 1%. It
was found that 1 mg of DHE, nasally administered, had the
equivalence of 10 mg orally, and almost 40% of the
bioavailability of the i.m. administration (PG Andersson
30 and LT Jespersen, Cephalalgia 1986; 6: 51-54).

- The maximal vasoconstrictor effect of 1 mg nasal DHE
amounted to about 40%, of 0.5 mg i.m. DHE to about 50% of
the initial venous diameter (W.H. Aellig and J. Rosenthaler,
35 Eur. J. Clin. Pharmacol. 1986; 30: 581-584).

Nasal DHE appeared to be equally effective than a
combination of oral ergotamine and caffeine in relieving

- migraine attacks (D. Kirt et al, Cephalalgia 1989; 9, suppl. 10: 410-411). Another study in 904 patients confirmed the efficacy of nasal DHE and reported side effects in 18.4% of patients: nasal irritation, nausea, vomiting, fatigue, vertigo, breathlessness, tachycardia and perspiration. Only 3.9% of the patients refused further treatment with nasal DHE (G. Jenzer and M.F. Bremgartner, Schweiz. Rundsch. Med. Prax. 1990; 79: 914-917). Lataste et al (Cephalalgia 1989; 9 suppl. 10: 342-343) and Di Serio et al (Cephalalgia 1989; 9 suppl. 10: 344-345), confirm the efficacy of nasal DHE in the acute management of migraine. In contrast, Tulunay et al (Cephalalgia 1987; 7: 131-133) found little difference in nasal DHE and placebo.
- Most of these studies are very encouraging and therefore nasal DHE, in the pharmaceutical composition studied by the above mentioned authors, seems an interesting alternative for oral and parenteral DHE preparations. Nasal DHE in the composition of DHE mesylate 4 mg/ml in 5% glucose and 1% caffeine, is available on prescription in several countries (e.g. Switzerland, France, Belgium).

Nevertheless, there is an urgent need for another DHE-nasal drug formulation, because the nasal preparation, presently on the market, is not stable. It is on the market as a separate glass ampoule (containing the DHE formulation) which has to be broken by the patient and sprayed in the nose using a separate spray device. After opening of the ampoule, the spray can be used no longer than 24 hours.

Accordingly, it is an object of the invention to provide a highly stable pharmaceutical composition, suitable for nasal administration, capable of introducing efficiently a therapeutical amount of DHE into the human body. It has surprisingly been found that a pharmaceutically acceptable DHE composition can be formulated, suitable for nasal administration, without the presence of a special

caffeine-glucose vehicle and without the necessity of presenting the formulation in a separate glass ampoule.

According to the invention, the nasal pharmaceutical composition contains DHE and/or a salt of DHE (mesylate or tartrate) and a cyclodextrin and/or other saccharides and/or sugar alcohols. Such compositions appear to result in a surprisingly high bioavailability and a superior stability of DHE.

10

The term "cyclodextrins" refers to cyclic oligosaccharides, like α -, β - and γ -cyclodextrin and their derivatives, preferably β -cyclodextrin and its derivatives, preferably methylated β -cyclodextrin, with a degree of CH_3 -substitution

15 between 0.5 and 3.0, more preferably between 1.7 and 2.1.

The term "saccharides" refers to disaccharides, like lactose, maltose, saccharose and also refers to polysaccharides, like dextrans, with an average molecular weight between 10.000 and 100.000, preferably 40.000 and

20 70.000. The term "sugar alcohols" refers to mannitol and sorbitol.

The nasal composition, according to the invention, can be administered as a nasal spray, nasal drop, suspension, gel, ointment, cream or powder. The administration of the nasal composition may also take place using a nasal tampon or nasal sponge, containing the invention composition.

In particular, powder formulations show a surprisingly high bioavailability and superior stability of the DHE. In addition, powder formulations have the advantage that no preservatives are necessary. Preservatives are known to decrease the ciliary movement, which may be harmful in chronic nasal medication (Hermens W.A.J.J. and Merkus F.W.H.M., Pharm. Res. 1987; 4: 445-449).

Nasal powder compositions can be made by mixing the active agent and the excipient, both possessing the desired particle size. Other methods to make a suitable powder formulation can be selected. Firstly, a solution of the active agent and the cyclodextrin and/or the other saccharide and/or sugar alcohol is made, followed by precipitation, filtration and pulverization. It is also possible to remove the solvent by freeze drying, followed by pulverization of the powder in the desired particle size by using conventional techniques, known from the pharmaceutical literature. The final step is size classification for instance by sieving, to get particles that are less than 100 microns in diameter, preferably between 50 and 100 microns in diameter. Powders can be administered using a nasal insufflator. Powders may also be administered in such a manner that they are placed in a capsule. The capsule is set in an inhalation or insufflation device. A needle is penetrated through the capsule to make pores at the top and the bottom of the capsule and air is sent to blow out the powder particles. Powder formulation can also be administered in a jet-spray of an inert gas or suspended in liquid organic fluids.

Also the active agent can be brought into a viscous basis, using vehicles, conventionally used, for example natural gums, methylcellulose and derivatives, acrylic polymers (carbopol) and vinyl polymers (polyvinylpyrrolidone). In the invention compositions many other excipients, known from the pharmaceutical literature, can be added, such as preservatives, surfactants, co-solvents, adhesives, anti-oxidants, buffers, viscosity enhancing agents, and agents to adjust the pH or the osmolarity.

The required amount for a nasal administration of a liquid or semi-solid nasal administration form is generally between 0.05 ml and 0.2 ml, preferably about 0.1 ml per nostril. The amount of a powder nasal formulation is

generally between 1 and 15 mg, preferably about 5 to 10 mg per nostril. Doses of DHE in the nasal pharmaceutical composition of the invention, suitable in the treatment of
5 migraine attacks, are preferably in the range from 0.25 to 0.5 mg per nostril.

The following examples illustrate the invention in more detail, but are not construed as limiting the invention:

10

EXAMPLE 1 (liquid)

Dihydroergotamine mesylate	250 mg
Methyl- β -cyclodextrin D.S. 1.8	2.5 g
Benzalkonium Chloride	0.01 %
15 Sodium EDTA	0.05-0.1 %
Sorbitol	5 %
Purified water to	100 ml
100 μ l = 250 μ g DHE mesylate	

20 EXAMPLE 2 (gel)

Dihydroergotamine mesylate	0.5 g
Methyl- β -cyclodextrin D.S. 1.8	5 g
Benzalkonium Chloride	0.01 %
Sodium EDTA	0.05-0.1 %
25 Sorbitol	5 %
Hydroxypropylmethylcellulose	1-2 %
Purified water to	100 ml
100 μ l gel = 500 μ g DHE	

30 EXAMPLE 3A (powder)

Dihydroergotamine mesylate	0.5 mg
Methyl- β -cyclodextrin	5 mg
Mannitol	4.5 mg
10 mg powder = 500 μ g DHE mesylate	

35

EXAMPLE 3B (powder)

Dihydroergotamine mesylate	0.5 mg
Dextran (average M.W. 70.000)	9.5 mg
10 mg powder = 500 µg DHE mesylate	

5

EXAMPLE 3C (powder)

Dihydroergotamine mesylate	0.5 mg
β-cyclodextrin	5 mg
Lactose	4.5 mg

10 10 mg powder = 500µg DHE mesylate

Apomorphine is a very potent dopamine agonist. It is used as an adjunctive medication in the treatment of Parkinson disease, complicated by motor fluctuations. Recently,

15 encouraging results have been reported on the intranasal application of apomorphine in patients with Parkinson disease to relieve "off-period" symptoms in patients with response fluctuations (T. van Laar et al, Arch. Neurol. 1992; 49: 482-484). The intranasal applied apomorphine,

20 used by these authors, consisted of an aqueous solution of apomorphine HCl 10 mg/ml. This formulation is also used for parenteral application and is published in different Pharmacopoeia's.

25 The exact nasal composition formulation used in the study by T. van Laar et al (1992) was:

Apomorphine HCl 0.5 H ₂ O	1 g
Sodium metabisulphite	0.100 g
30 Sodium EDTA	0.010 g
NaCl	0.600 g
Benzalkonium Chloride	0.01 %
NaH ₂ PO ₄ ·2H ₂ O	0.150 g
Na ₂ HPO ₄ ·2H ₂ O	0.050 g
35 NaOH 1 M to adjust pH at 5.8	
purified water to 100 ml	

(from Pharm. Weekblad 1991; 126: 1113-1114)

By a metered dose nebulizer a dose of 1 mg apomorphine HCl (0.1 ml of the solution) was delivered with each nasal application by puff to the patients. A great disadvantage of this aqueous solution is the instability of the apomorphine.

An object of the invention is a nasal formulation of apomorphine with an improved bioavailability and stability of apomorphine.

10

According to the invention, the nasal pharmaceutical composition contains apomorphine and/or apomorphine salts and a cyclodextrin and/or other saccharides and/or sugar alcohols. Such compositions appear to result in a surprisingly high bioavailability and superior stability of apomorphine.

The term "cyclodextrins" refers to cyclic oligosaccharides, like α -, β - and γ -cyclodextrin and their derivatives, preferably β -cyclodextrin and its derivatives, preferably methylated β -cyclodextrin, with a degree of CH_3 -substitution between 0.5 and 3.0, more preferably between 1.7 and 2.1. The term "saccharides" refers to disaccharides, like lactose, maltose, saccharose and also refers to polysaccharides, like dextrans, with an average molecular weight between 10.000 and 100.000, preferably 40.000 and 70.000. The term "sugar alcohols" refers to mannitol and sorbitol.

The nasal composition, according to the invention, can be administered as a nasal spray, nasal drop, suspension, gel, ointment, cream or powder. The administration of the nasal composition may also take place using a nasal tampon or nasal sponge, containing the invention composition.

35

In particular, powder formulations show a surprisingly high bioavailability and superior stability of the apomorphine.

In addition, powder formulations have the advantage that no preservatives are necessary. Preservatives are known to decrease the ciliary movement, which may be harmful in chronic nasal medication (Hermens W.A.J.J. and Merkus F.W.H.M., Pharm. Res. 1987; 4: 445-449).

Nasal powder compositions can be made by mixing the active agent and the excipient, both possessing the desired particle size. Other methods to make a suitable powder formulation can be selected. Firstly, a solution of the active agent and the cyclodextrin and/or the other saccharide and/or sugar alcohol is made, followed by precipitation, filtration and pulverization. It is also possible to remove the solvent by freeze drying, followed by pulverization of the powder in the desired particle size by using conventional techniques, known from the pharmaceutical literature. The final step is size classification for instance by sieving, to get particles that are less than 100 microns in diameter, preferably between 50 and 100 microns in diameter. Powders can be administered using a nasal insufflator. Powders may also be administered in such a manner that they are placed in a capsule. The capsule is set in an inhalation or insufflation device. A needle is penetrated through the capsule to make pores at the top and the bottom of the capsule and air is sent to blow out the powder particles. Powder formulation can also be administered in a jet-spray of an inert gas or suspended in liquid organic fluids.

Also the active agent can be brought into a viscous basis, using vehicles, conventionally used, for example natural gums, methylcellulose and derivatives, acrylic polymers (carbopol) and vinyl polymers (polyvinylpyrrolidone). In the invention compositions many other excipients, known from the pharmaceutical literature, can be added, such as preservatives, surfactants, co-solvents, adhesives,

anti-oxidants, buffers, viscosity enhancing agents, and agents to adjust the pH or the osmolarity.

The required amount for a nasal administration of a liquid or semi-solid nasal administration form is generally between 0.05 ml and 0.2 ml, preferably about 0.1 ml per nostril. The amount of a powder nasal formulation is generally between 1 and 15 mg, preferably about 5 to 10 mg per nostril. Doses of apomorphine in the pharmaceutical composition of the present invention, suitable in the treatment of Parkinson disease, are generally in the range of 0.1 to 2mg, preferably between 0.5 mg and 1 mg per nostril.

The following examples illustrate the present invention in more detail, but are not construed as limiting the invention:

EXAMPLE 1A (powder)

20	Apomorphine base	1 mg
	Methyl- β -cyclodextrin D.S. 2.1	5 mg
	Mannitol	4 mg
	10 mg powder = 1 mg Apomorphine	

25 EXAMPLE 1B (powder)

	Apomorphine HCl	2 mg
	Mannitol	18 mg
	20 mg powder = 2 mg Apomorphine HCl	

30 EXAMPLE 1C (powder)

	Apomorphine HCl	1 mg
	Dextran (average M.W. 70.000)	9 mg
	10 mg powder = 1 mg Apomorphine HCl	

EXAMPLE 2 (gel)

	Apomorphine HCl	500 mg
	Methylated- β -cyclodextrin D.S. 1.8	2.5 g
	Hydroxypropylmethylcellulose	1-2 g
5	Benzalkonium Chloride	0.01 %
	Sodium EDTA	0.1 %
	Sodium metabisulphite	0.15 %
	Sorbitol	4 %
	pH adjusted to	4.5 - 5.5
10	purified water to	100 ml
	0.2 ml gel = 1 mg Apomorphine HCl	

EXAMPLE 3 (liquid)

	Apomorphine HCl	1 g
15	Methylated- β -cyclodextrin D.S. 1.1	4 g
	Sodium metabisulphite	0.15 %
	Sodium EDTA	0.1 %
	Benzalkonium Chloride	0.01 %
	NaCl	0.8 %
20	pH adjusted to	4.5 - 5.5
	purified water to	100 ml
	100 μ l = 1 mg Apomorphine HCl	

- Morphine is one of the strongest analgesics. Morphine
- 25 therapy is restricted to two groups of patients. Firstly, to hospitalized patients, after surgery and secondly, to cancer and burn patients. The latter treatment is chronic. Morphine is administered generally by injection and in chronic treatment by sustained release oral preparations.
- 30 After single oral administration morphine has a poor effect, mainly due to a large first pass effect. Secondly, the oral route is not possible when the patient shows severe nausea, vomiting, bowel obstruction or confusion. There is a need for a non-parenteral administration, other
- 35 then oral, because injection therapy needs interference of (para)medical personnel and is painful.

Buccal administration of morphine have been proposed (MDD Bell et al, Lancet 1985; 1: 71-73), but this route did not find a large acceptance in practice. Recently rectal administration of morphine has been studied (T.J. Wilkinson et al, Cancer Chemother. Pharmacol 1992; 31: 251-254 and N Babul et al Clin. Pharmacol. Ther. 1993; 54: 286-292). From both publications it can be concluded that rectal application in some cases may be an alternative when the parenteral route is impractical or undesirable and the oral route is not available due to the patients condition. Nasal administration of a strong analgesic could be a good alternative to parenteral therapy, because it may give a very rapid absorption and no first pass effect.

To overcome the drawbacks of the oral and parenteral routes of administration of morphine, the use of a nasal spray has been proposed (S.L. Verweij and R. van Gijn: Can morphine be administered nasally? Ziekenhuisfarmacie (Dutch) 1988; 4: 73-77). The composition of the nasal spray in this study was:

Morphine HCl.3H ₂ O	1.50 g
Sodium metabisulphite	0.03 g
Sodium EDTA	0.003 g
25 Benzylalcohol	0.3 ml
Propylene glycol	6 ml
Phosphate Buffer (0.01 mol/L; pH 6.00)	30 ml
Per puff of 100 µl the dose of morphine is	5 mg.

In 7 volunteers Verweij and van Gijn studied the pharmacokinetics of morphine after 4 puffs of about 100 µl (2 times 1 puff of 100, µl in each nostril). The exact dose which was delivered to the volunteers was 16 mg of morphine (range 15-18 mg) and the bioavailability of morphine from this nasal spray was 26-35%. The bioavailability of morphine after oral application is estimated to be about 40% (J. Säwe, Clin. Pharmacokinetics

1986; 11: 87-106). This means, that the bioavailability of morphine after giving the nasal spray as described by Verweij and van Gijn is relatively low. After nasal absorption there is no first pass effect and therefore the
5 nasal bioavailability should be higher than the oral.

The nasal absorption of morphine has been studied also by F Chast et al (J. Pharm. Clin. 1992; 11: 257-261). They delivered nasally and orally 20 mg morphine acetate in an
10 aqueous solution to 6 patients and compared the nasal absorption with the oral absorption of the same solution. They found, as expected, higher blood levels of morphine after the nasal application. Unfortunately, the nasal
15 solutions, as described by the preceding studies of Verweij and van Gijn and of Chast and coworkers, are not stable and the bioavailability of morphine can be improved.

An object of the invention is to provide a highly stable pharmaceutical composition, suitable for nasal
20 administration, and showing an superior bioavailability of morphine.

According to the invention, the nasal pharmaceutical composition contains morphine and/or morphine salts
25 (hydrochloride, sulphate, acetate) and a cyclodextrin and/or other saccharides and/or sugar alcohols. Such compositions appear to result in a surprisingly high bioavailability and superior stability of morphine.

30 The term "cyclodextrins" refers to cyclic oligosaccharides, like α -, β - and γ -cyclodextrin and their derivatives, preferably β -cyclodextrin and its derivatives, preferably methylated β -cyclodextrin, with a degree of CH_3 -substitution between 0.5 and 3.0, more preferably between 1.7 and 2.1.
35 The term "saccharides" refers to disaccharides, like lactose, maltose, saccharose and also refers to polysaccharides, like dextrans, with an average molecular

weight between 10.000 and 100.000, preferably 40.000 and 70.000. The term "sugar alcohols" refers to mannitol and sorbitol.

- 5 The nasal composition, according to the invention, can be administered as a nasal spray, nasal drop, suspension, gel, ointment, cream or powder. The administration of the nasal composition may also take place using a nasal tampon or nasal sponge, containing the invention composition.

10

- In particular, powder formulations show a surprisingly high bioavailability and superior stability of the morphine. In addition, powder formulations have the advantage that no preservatives are necessary. Preservatives are known to
15 decrease the ciliary movement, which may be harmful in chronic nasal medication (Hermens W.A.J.J. and Merkus F.W.H.M., Pharm. Res. 1987; 4: 445-449).

- Nasal powder compositions can be made by mixing the active
20 agent and the excipient, both possessing the desired particle size. Other methods to make a suitable powder formulation can be selected. Firstly, a solution of the active agent and the cyclodextrin and/or the other
saccharide and/or sugar alcohol is made, followed by
25 precipitation, filtration and pulverization. It is also possible to remove the solvent by freeze drying, followed by pulverization of the powder in the desired particle size by using conventional techniques, known from the pharmaceutical literature. The final step is size
30 classification for instance by sieving, to get particles that are less than 100 microns in diameter, preferably between 50 and 100 microns in diameter. Powders can be administered using a nasal insufflator. Powders may also be administered in such a manner that they are placed in a
35 capsule. The capsule is set in an inhalation or insufflation device. A needle is penetrated through the capsule to make pores at the top and the bottom of the

capsule and air is sent to blow out the powder particles. Powder formulation can also be administered in a jet-spray of an inert gas or suspended in liquid organic fluids.

- 5 Also the active agent can be brought into a viscous basis, using vehicles, conventionally used, for example natural gums, methylcellulose and derivatives, acrylic polymers (carbopol) and vinyl polymers (polyvinylpyrrolidone). In the invention compositions many other excipients, known
10 from the pharmaceutical literature, can be added, such as preservatives, surfactants, co-solvents, adhesives, anti-oxidants, buffers, viscosity enhancing agents, and agents to adjust the pH or the osmolarity.
- 15 The required amount for a nasal administration of a liquid or semi-solid nasal administration form is generally between 0.05 ml and 0.2 ml, preferably about 0.1 ml per nostril. The amount of a powder nasal formulation is generally between 1 and 15 mg, preferably about 5 to 10 mg
20 per nostril.

Doses of morphine in the pharmaceutical composition of the present invention, suitable in the treatment of pain, are in the range from 1 to 20 mg.

25

The following examples illustrate the present invention in more detail, but are not construed as limiting the invention:

30 EXAMPLE 1A (powder)

Morphine sulphate 5H ₂ O	13.3 mg
Methyl- β -cyclodextrin D.S. 2.1	11.7 mg
Mannitol	5 mg

30 mg powder = 10 mg morphine

35

EXAMPLE 1B (powder)

Morphine sulphate 5H ₂ O	13.3 mg
β-cyclodextrin	6.7 mg
20 mg powder = 10 mg morphine	

5

EXAMPLE 1C (powder)

Morphine HCl 3H ₂ O	13.1 mg
Dextran (average MW 70.000)	16.9 mg
30 mg powder = 10 mg morphine	

10

EXAMPLE 2 (gel)

Morphine (as salt)	1.5 g
Methyl-β-cyclodextrin D.S. 1.8	5 g
(Hydroxypropyl)methylcellulose	1-2 %
15 Benzalkonium Chloride	0.01 %
Sodium EDTA	0.1 %
Sodium metabisulphite	0.15 %
Sorbitol	4 %
Purified water to	50 ml
20 0.2 ml gel = 6 mg morphine	

EXAMPLE 3 (liquid)

Morphine (as salt)	4 g
Methyl-β-cyclodextrin D.S. 2.1	4 g
25 Methylcellulose	0.25 %
Sodium metabisulphite	0.15 %
Sodium EDTA	0.1 %
Benzalkonium Chloride	0.01 %
Mannitol	4 %
30 Purified water to	100 ml
100 μl = 4 mg morphine	

CLAIMS

1. A pharmaceutical composition for nasal administration comprising dihydroergotamine and/or a
5 dihydroergotamine-salt and a cyclodextrin and/or an other saccharide and/or a sugar alcohol.
2. A pharmaceutical composition according to claim 1,
10 wherein the dihydroergotamine salt is dihydroergotamine mesylate and/or tartrate.
3. A pharmaceutical composition according to claim 1 or 2,
15 wherein the cyclodextrin is α - or β - or γ -cyclodextrin and/or a derivative of α - or β - or γ -cyclodextrin.
4. A pharmaceutical composition according to any of claims
1 - 3, wherein the cyclodextrin is β -cyclodextrin and/or a derivative.
- 20 5. A pharmaceutical composition according to claim 4,
wherein the derivative of β -cyclodextrin is a methylated β -cyclodextrin with a degree of substitution between 0.5 and 3.0.
- 25 6. A pharmaceutical composition according to any of claims
1 - 5, wherein the saccharides a disaccharide, preferably lactose and/or a polysaccharide, preferably dextran, having an average molecular weight between
30 10.000 and 100.000, preferably between 40.000 and 70.000 .
7. A pharmaceutical composition according to any of claims
35 1-6, wherein the sugar alcohol is mannitol and/or sorbitol.

8. A pharmaceutical composition according to any of claims 1-7 in powder form, suitable for nasal administration, wherein the particles of the powder have a diameter between 50-100 microns.
- 5
9. A process of preparing a nasal pharmaceutical composition according to any of claims 1 - 8, which process comprises combining dihydroergotamine and/or its salts with a cyclodextrin and/or a disaccharide and/or a polysaccharide and/or a sugar alcohol.
- 10
10. A method of treating migraine attacks by administering a pharmaceutical composition, according to any of claims 1 - 8, to the nasal mucosa.
- 15
11. A pharmaceutical composition for nasal administration comprising apomorphine and/or an apomorphine salt and a cyclodextrin and/or an other saccharide and/or a sugar alcohol.
- 20
12. A pharmaceutical composition according to claim 11, wherein the apomorphine salt is apomorphine hydrochloride.
- 25
13. A pharmaceutical composition according to claim 11 or 12, wherein the cyclodextrin is α - or β - or γ -cyclodextrin and/or a derivative of α - or β - or γ -cyclodextrin.
- 30
14. A pharmaceutical composition according to any of claims 11 - 13, wherein the cyclodextrin is β -cyclodextrin and/or a derivative.

15. A pharmaceutical composition according to claim 14,
wherein the derivative of β -cyclodextrin is a
methyalted β -cyclodextrin with a degree of substitution
5 between 0.5 and 3.0.
16. A pharmaceutical composition according to any of claims
11 - 15, wherein the saccharide is a disaccharide,
preferably lactose and/or a polysaccharide, preferably
10 dextran, having an average molecular weight between
10.000 and 100.000, preferably between 40.000 and
70.000.
17. A pharmaceutical composition according to any of claims
11 - 16, wherein the sugar alcohol is mannitol and/or
15 sorbitol.
18. A pharmaceutical composition according to any of claims
11 - 17 in powder form, suitable for nasal
20 administration, wherein the particles of the powder
have a diameter between 50-100 microns.
19. A process of preparing a nasal pharmaceutical
composition according to any of claims 11 - 18, which
25 process comprises combining apomorphine and/or its
salts with a cyclodextrin and/or a disaccharide and/or
a polysaccharide and/or a sugar alcohol.
20. A method of treating Parkinson disease by administering
30 a pharmaceutical composition according to any of claims
11 - 18, to the nasal mucosa.

21. A pharmaceutical composition for nasal administration comprising morphine and/or morphine salts and a cyclodextrin and/or an other saccharide and/or a sugar alcohol.
22. A pharmaceutical composition according to claim 21, wherein the morphine salt is morphine hydrochloride and/or acetate and/or sulphate.
23. A pharmaceutical composition according to claim 21 or 22, wherein the cyclodextrin is α - or β - or γ -cyclodextrin and/or a derivative of α - or β - or γ -cyclodextrin.
24. A pharmaceutical composition according to any of claims 21 - 23, wherein the cyclodextrin is β -cyclodextrin and/or a derivative.
25. A pharmaceutical composition according to claim 24, wherein the derivative of β -cyclodextrin is a methylated β -cyclodextrin with a degree of substitution between 0.5 and 3.0.
26. A pharmaceutical composition according to any of claims 21 - 25, wherein the saccharide is a disaccharide, preferably lactose and/or a polysaccharide, preferably dextran, having an average molecular weight between 10.000 and 100.000, preferably between 40.000 and 70.000.
27. A pharmaceutical composition according to any of claims 21 - 26, wherein the sugar alcohol is mannitol and/or sorbitol.

28. A pharmaceutical composition according to any of claims
21 - 27 in powder form, suitable for nasal
administration, wherein the particles of the powder
5 have a diameter between 50-100 microns.
29. A process of preparing a nasal pharmaceutical
composition, according to any of claims 21 - 28, which
• process comprises combining morphine and/or its salts
10 with a cyclodextrin and/or a disaccharide and/or a
polysaccharide and/or a sugar alcohol.
30. A method of relieving pain by administering a
pharmaceutical composition according to any of claims
15 21 - 28, to the nasal mucosa.